


CLEAN VERSION OF AMENDED CLAIMS

1. (Twice Amended) A molecular compound selected from the group consisting of hydrates, solvates, adducts and clathrate compounds prepared by the method of reacting a phenol derivative represented by Formula (I)





wherein R_1 and R_5 are same or different selected from the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons, hydroxyl or



wherein Y and Z are selected from the group consisting of alkyl having 1 to 8 carbons, alkenyl having 2 to 8 carbons, alkoxy having 1 to 6 carbons, hydroxyl, substituted amino, substituted cycloalkyl, substituted phenyl or substituted aralkyl;



R_2 and R_4 are same or different selected from the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons or

hydroxyl or




wherein Y and Z are as defined above, in case R_1 , R_3 or R_5 is alkoxy having 1 to 4 carbons or hydroxyl;

R_3 is selected from the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons, hydroxyl, Formula (II) or Formula (III)




wherein X is selected from the group consisting of



wherein w is 0, 1 or 2; u is 0 or 1; q is 0 to 4; R₁₄ and R₁₅ are same or different selected from the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons, hydroxyl, optionally substituted phenyl or optionally substituted aralkyl; R₁₆ is selected from the group consisting of hydrogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons, hydroxyl, substituted phenyl or substituted aralkyl;

R₆, R₉ and R₁₀ are same or different selected from the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons, hydroxyl, or



wherein Y and Z are as defined above;

R₇, R₈, R₁₁ and R₁₃ are same or different selected from the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons or hydroxyl, but R₁₁ is selected from the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons, hydroxyl or



wherein Y and Z are as defined above in case R₁₂ is alkoxy having 1 to 4 carbons or hydroxyl;

R₁₂ is selected from the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons, hydroxyl or selected from the group consisting of



wherein Y and Z are as defined above, or selected from the group consisting of



21,

wherein Y and Z are as defined above, or

when R₃ is of Formula (II), one of R₁, R₅, R₆ and R₉ is selected from the group consisting of



wherein Y and Z are as defined above


when R₃ is of Formula (III), at least one of R₁, R₅ and R₁₀ is selected from the group consisting of



where Y and Z are as defined above, and

when R₃ is selected from a group other than the group consisting of Formula (II) or (III), either R₁ or R₅ is selected from the group consisting of


201



wherein Y and Z are as defined above, and


the phenol derivative is reacted with an organic compound under conditions sufficient to form the molecular compound selected from the group consisting of hydrates, solvates, adducts and clathrate compounds having the phenol derivative as a constituent, the constituent being a host.

2. (Twice Amended) A molecular compound selected from the group consisting of hydrates, solvates, adducts and clathrate compounds prepared by the method of reacting a phenol derivative represented by Formula (IV)



wherein A is selected from the group consisting of

201



wherein w is 0, 1 or 2 and u is 0 or 1; R₁₈, R₁₉, R₂₁ and R₂₄ are same or different selected from the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons; R₁₇ is selected from the group consisting of




wherein Y and Z are selected from the group consisting of alkyl having 1 to 6 carbons, alkenyl

201


having 2 to 6 carbons, cyclohexyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, cyclopentyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, phenyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or halogen, benzyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, phenethyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, α -methylbenzyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, or naphthyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, and R_{20} , R_{22} and R_{23} are same or different, hydrogen, halogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons or the same groups as those for R_{17} , and

an organic compound, as the other reactant under conditions sufficient to form the molecular compound selected from the group consisting of hydrates, solvates, adducts and clathrate compounds having the phenol derivative as a constituent, the constituent being a host.

3. (Amended) A molecular compound selected from the group consisting of hydrates, solvates, adducts and clathrate compounds prepared by the method of reacting a phenol derivative represented by Formula (V)



wherein B is a group selected from




wherein w is 0, 1 or 2 and u is 0 or 1; R₂₆, R₂₇, R₃₀ and R₃₂ are same or different selected from

the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons; R_{25} , R_{28} , R_{29} and R_{31} are same or different selected from the group consisting of hydrogen, halogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4 carbons or




wherein Y and Z are selected from the group consisting of alkyl having 1 to 6 carbons, alkenyl having 2 to 6 carbons, cyclohexyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, cyclopentyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, phenyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or halogen, benzyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, phenethyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, α -methylbenzyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, or naphthyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, and at least one of R_{25} , R_{28} and R_{29} is selected from the group consisting of



wherein Y and Z are selected from the group consisting of alkyl having 1 to 6 carbons, alkenyl having 2 to 6 carbons, cyclohexyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, cyclopentyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, phenyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or halogen, benzyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, phenethyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, α -methylbenzyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, or naphthyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, and an organic compound as the second reactant under conditions sufficient to form the molecular compound selected from the group consisting of hydrates, solvates, adducts and clathrate compounds having the phenol derivative as a constituent, the constituent being a host.

4. (Amended) A molecular compound selected from the group consisting of hydrates, solvates, adducts and clathrate compounds prepared by the method of reacting a phenol derivative represented by Formula (VI)



wherein R_{33} is selected from the group consisting of



wherein Y and Z are selected from the group consisting of alkyl having 1 to 6 carbons, alkenyl having 2 to 6 carbons, cyclohexyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, cyclopentyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, phenyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or halogen, benzyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or halogen, phenethyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons

or alkoxy having 1 to 4 carbons or hydroxyl or halogen,
 α -methylbenzyl which may have alkyl having 1 to 4 carbons or alkenyl having 2 to 4 carbons or
alkoxy having 1 to 4 carbons or hydroxyl or halogen, or naphthyl which may have alkyl having 1
to 4 carbons or alkenyl having 2 to 4 carbons or alkoxy having 1 to 4 carbons or hydroxyl or
halogen, and R_{34} , R_{35} , R_{36} and R_{37} are same or different selected from the group consisting of
hydrogen, alkyl having 1 to 4 carbons, alkenyl having 2 to 4 carbons, alkoxy having 1 to 4
carbons, hydroxyl, halogen or the same groups as those for R_{33} , with an organic compound as the
second reactant under conditions sufficient to form the molecular compound selected from the
group consisting of hydrates, solvates, adducts and clathrate compounds having the phenol
derivative as a constituent, the constituent being a host.

NEW CLAIMS

28. The molecular compound prepared according to the method of claim 1, wherein the organic compound is selected from the group comprising:

antibacterial agents, antifungal agents, insecticides, noxious insect repellants, perfumes, deodorants, antifouling agents, curing agents and accelerators for coating materials, resins and adhesives, natural essential oils, antioxidants, vulcanization accelerators or organic solvents, that react with the said phenol derivative to form the molecular compound.

29. The molecular compound prepared according to the method of claim 2, wherein the organic compound is selected from the group comprising:

antibacterial agents, antifungal agents, insecticides, noxious insect repellants, perfumes, deodorants, antifouling agents, curing agents and accelerators for coating materials, resins and adhesives, natural essential oils, antioxidants, vulcanization accelerators or organic solvents, that react with the said phenol derivative to form the molecular compound.


30. The molecular compound prepared according to the method of claim 3, wherein the organic compound is selected from the group comprising:

antibacterial agents, antifungal agents, insecticides, noxious insect repellants, perfumes, deodorants, antifouling agents, curing agents and accelerators for coating materials, resins and adhesives, natural essential oils, antioxidants, vulcanization accelerators or organic solvents, that react with the said phenol derivative to form the molecular compound.

102

31. The molecular compound prepared according to the method of claim 4, wherein the organic compound is selected from the group comprising:

antibacterial agents, antifungal agents, insecticides, noxious insect repellants, perfumes, deodorants, antifouling agents, curing agents and accelerators for coating materials, resins and adhesives, natural essential oils, antioxidants, vulcanization accelerators or organic solvents, that react with the said phenol derivative to form the molecular compound.

A handwritten signature, possibly 'JL', is located on the left side of the page. A long, slightly curved horizontal line extends from the signature across the middle of the page.